

A GENERALISED TRAPEZOID TYPE INEQUALITY FOR CONVEX FUNCTIONS

S.S. DRAGOMIR

ABSTRACT. A generalised trapezoid inequality for convex functions and applications for quadrature rules are given. A refinement and a counterpart result for the Hermite-Hadamard inequalities are obtained and some inequalities for pdf's and (HH) -divergence measure are also mentioned.

1. INTRODUCTION

The following integral inequality for the generalised trapezoid formula was obtained in [2] (see also [1, p. 68]):

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation. We have the inequality*

$$(1.1) \quad \left| \int_a^b f(t) dt - [(x-a)f(a) + (b-x)f(b)] \right| \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(f),$$

holding for all $x \in [a, b]$, where $\bigvee_a^b(f)$ denotes the total variation of f on the interval $[a, b]$.

The constant $\frac{1}{2}$ is the best possible one.

This result may be improved if one assumes the monotonicity of f as follows (see [1, p. 76])

Theorem 2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a monotonic nondecreasing function on $[a, b]$. Then we have the inequality:*

$$(1.2) \quad \begin{aligned} & \left| \int_a^b f(t) dt - [(x-a)f(a) + (b-x)f(b)] \right| \\ & \leq (b-x)f(b) - (x-a)f(a) + \int_a^b \operatorname{sgn}(x-t)f(t) dt \\ & \leq (x-a)[f(x) - f(a)] + (b-x)[f(b) - f(x)] \\ & \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] [f(b) - f(a)] \end{aligned}$$

Date: June 26, 2001.

1991 Mathematics Subject Classification. Primary 26D14; Secondary 26D99.

Key words and phrases. Generalised Trapezoid Inequality, Hermite-Hadamard Inequality, Probability density functions, Divergence measures.

for all $x \in [a, b]$.

The above inequalities are sharp.

If the mapping is Lipschitzian, then the following result holds as well [3] (see also [1, p. 82]).

Theorem 3. Let $f : [a, b] \rightarrow \mathbb{R}$ be an L -Lipschitzian function on $[a, b]$, i.e., f satisfies the condition:

$$(L) \quad |f(s) - f(t)| \leq L |s - t| \quad \text{for any } s, t \in [a, b] \quad (L > 0 \text{ is given}).$$

Then we have the inequality:

$$(1.3) \quad \left| \int_a^b f(t) dt - [(x-a)f(a) + (b-x)f(b)] \right| \leq \left[\frac{1}{4}(b-a)^2 + \left(x - \frac{a+b}{2}\right)^2 \right] L$$

for any $x \in [a, b]$.

The constant $\frac{1}{4}$ is best in (1.3).

If we would assume absolute continuity for the function f , then the following estimates in terms of the Lebesgue norms of the derivative f' hold [1, p. 93].

Theorem 4. Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$. Then for any $x \in [a, b]$, we have

$$(1.4) \quad \left| \int_a^b f(t) dt - [(x-a)f(a) + (b-x)f(b)] \right| \leq \begin{cases} \left[\frac{1}{4}(b-a)^2 + \left(x - \frac{a+b}{2}\right)^2 \right] \|f'\|_\infty & \text{if } f' \in L_\infty[a, b]; \\ \frac{1}{(q+1)^{\frac{1}{q}}} \left[(x-a)^{q+1} + (b-x)^{q+1} \right]^{\frac{1}{q}} \|f'\|_p & \text{if } f' \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[\frac{1}{2}(b-a) + \left|x - \frac{a+b}{2}\right| \right] \|f'\|_1, & \end{cases}$$

where $\|\cdot\|_p$ ($p \in [1, \infty]$) are the Lebesgue norms, i.e.,

$$\|f'\|_\infty = \operatorname{ess\,sup}_{s \in [a, b]} |f'(s)|$$

and

$$\|f'\|_p := \left(\int_a^b |f'(s)|^p ds \right)^{\frac{1}{p}}, \quad p \geq 1.$$

In this paper we point out some similar results for convex functions. Applications for quadrature formulae, for probability density functions and HH -Divergences in Information Theory are also considered.

2. THE RESULTS

The following theorem providing a lower bound for the difference

$$(x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt$$

holds.

Theorem 5. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$. Then for any $x \in (a, b)$ we have the inequality*

$$(2.1) \quad \begin{aligned} & \frac{1}{2} \left[(b-x)^2 f'_+(x) - (x-a)^2 f'_-(x) \right] \\ & \leq (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt. \end{aligned}$$

The constant $\frac{1}{2}$ in the left hand side of (2.1) is sharp in the sense that it cannot be replaced by a larger constant.

Proof. It is easy to see that for any locally absolutely continuous function $f : (a, b) \rightarrow \mathbb{R}$, we have the identity

$$(2.2) \quad (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt = \int_a^b (t-x)f'(t) dt$$

for any $x \in (a, b)$, where f' is the derivative of f which exists a.e. on $[a, b]$.

Since f is convex, then it is locally Lipschitzian and thus (2.2) holds. Moreover, for any $x \in (a, b)$, we have the inequalities:

$$(2.3) \quad f'(t) \leq f'_-(x) \quad \text{for a.e. } t \in [a, x]$$

and

$$(2.4) \quad f'(t) \geq f'_+(x) \quad \text{for a.e. } t \in [x, b].$$

If we multiply (2.3) by $x-t \geq 0$, $t \in [a, x]$ and integrate on $[a, x]$, we get

$$(2.5) \quad \int_a^x (x-t)f'(t) dt \leq \frac{1}{2}(x-a)^2 f'_-(x)$$

and if we multiply (2.4) by $t-x \geq 0$, $t \in [x, b]$ and integrate on $[x, b]$, we also have

$$(2.6) \quad \int_x^b (t-x)f'(t) dt \geq \frac{1}{2}(b-x)^2 f'_+(x).$$

Finally, if we subtract (2.5) from (2.6) and use the representation (2.2), we deduce the desired inequality (2.1).

Now, assume that (2.1) holds with a constant $C > 0$ instead of $\frac{1}{2}$, i.e.,

$$(2.7) \quad \begin{aligned} & C \left[(b-x)^2 f'_+(x) - (x-a)^2 f'_-(x) \right] \\ & \leq (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt. \end{aligned}$$

Consider the convex function $f_0(t) := k \left| t - \frac{a+b}{2} \right|$, $k > 0$, $t \in [a, b]$. Then

$$\begin{aligned} f'_{0+} \left(\frac{a+b}{2} \right) &= k, \quad f'_{0-} \left(\frac{a+b}{2} \right) = -k, \\ f_0(a) &= \frac{k(b-a)}{2} = f_0(b), \quad \int_a^b f_0(t) dt = \frac{1}{4}k(b-a)^2. \end{aligned}$$

If in (2.7) we choose f_0 as above and $x = \frac{a+b}{2}$, then we get

$$C \left[\frac{1}{4}(b-a)^2 k + \frac{1}{4}(b-a)^2 k \right] \leq \frac{1}{4}k(b-a)^2$$

giving $C \leq \frac{1}{2}$, and the sharpness of the constant is proved. ■

Now, recall that the following inequality which is well known in the literature as the *Hermite-Hadamard inequality* for convex functions holds

$$(H-H) \quad f \left(\frac{a+b}{2} \right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a) + f(b)}{2}.$$

The following corollary gives a sharp lower bound for the difference

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt.$$

Corollary 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$. Then*

$$\begin{aligned} (2.8) \quad 0 &\leq \frac{1}{8} \left[f'_+ \left(\frac{a+b}{2} \right) - f'_- \left(\frac{a+b}{2} \right) \right] (b-a) \\ &\leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt. \end{aligned}$$

The constant $\frac{1}{8}$ is sharp.

The proof is obvious by the above theorem. The sharpness of the constant is obtained for $f_0(t) = k \left| t - \frac{a+b}{2} \right|$, $t \in [a, b]$, $k > 0$.

When x is a point of differentiability, we may state the following corollary as well.

Corollary 2. *Let f be as in Theorem 5. If $x \in (a, b)$ is a point of differentiability for f , then*

$$(2.9) \quad (b-a) \left(\frac{a+b}{2} - x \right) f'(x) \leq (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt.$$

Remark 1. *If $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is convex on I and if we choose $x \in \overset{\circ}{I}$ ($\overset{\circ}{I}$ is the interior of I), $b = x + \frac{h}{2}$, $a = x - \frac{h}{2}$, $h > 0$ is such that $a, b \in I$, then from (2.1) we may write*

$$(2.10) \quad 0 \leq \frac{1}{8}h^2 [f'_+(x) - f'_-(x)] \leq \frac{f(a) + f(b)}{2} \cdot h - \int_{x-\frac{h}{2}}^{x+\frac{h}{2}} f(t) dt$$

and the constant $\frac{1}{8}$ is sharp in (2.10).

The following result providing an upper bound for the difference

$$(x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt$$

also holds.

Theorem 6. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$. Then for any $x \in [a, b]$, we have the inequality:*

$$(2.11) \quad \begin{aligned} & (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt \\ & \leq \frac{1}{2} \left[(b-x)^2 f'_-(b) - (x-a)^2 f'_+(a) \right]. \end{aligned}$$

The constant $\frac{1}{2}$ is sharp in the sense that it cannot be replaced by a smaller constant.

Proof. If either $f'_+(a) = -\infty$ or $f'_-(b) = +\infty$, then the inequality (2.11) evidently holds true.

Assume that $f'_+(a)$ and $f'_-(b)$ are finite.

Since f is convex on $[a, b]$, we have

$$(2.12) \quad f'(t) \geq f'_+(a) \quad \text{for a.e. } t \in [a, x]$$

and

$$(2.13) \quad f'(t) \leq f'_-(b) \quad \text{for a.e. } t \in [x, b].$$

If we multiply (2.12) by $(x-t) \geq 0$, $t \in [a, x]$ and integrate on $[a, x]$, then we deduce

$$(2.14) \quad \int_a^x (x-t) f'(t) dt \geq \frac{1}{2} (x-a)^2 f'_+(a)$$

and if we multiply (2.13) by $t-x \geq 0$, $t \in [x, b]$ and integrate on $[x, b]$, then we also have

$$(2.15) \quad \int_x^b (t-x) f'(t) dt \leq \frac{1}{2} (b-x)^2 f'_-(b).$$

Finally, if we subtract (2.14) from (2.15) and use the representation (2.2), we deduce the desired inequality (2.11).

Now, assume that (2.11) holds with a constant $D > 0$ instead of $\frac{1}{2}$, i.e.,

$$(2.16) \quad \begin{aligned} & (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt \\ & \leq D \left[(b-x)^2 f'_-(b) - (x-a)^2 f'_+(a) \right]. \end{aligned}$$

If we consider the convex function $f_0 : [a, b] \rightarrow \mathbb{R}$, $f_0(t) = k|t - \frac{a+b}{2}|$, then we have $f'_-(b) = k$, $f'_+(a) = -k$ and by (2.16) we deduce for $x = \frac{a+b}{2}$ that

$$\frac{1}{4}k(b-a)^2 \leq D \left[\frac{1}{4}k(b-a)^2 + \frac{1}{4}k(b-a)^2 \right]$$

giving $D \geq \frac{1}{2}$, and the sharpness of the constant is proved. ■

The following corollary related to the Hermite-Hadamard inequality is interesting as well.

Corollary 3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be convex on $[a, b]$. Then*

$$(2.17) \quad 0 \leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{1}{8} [f'_-(b) - f'_+(a)] (b-a)$$

and the constant $\frac{1}{8}$ is sharp.

Remark 2. Denote $B := f'_-(b)$, $A := f'_+(a)$ and assume that $B \neq A$, i.e., f is not constant on (a, b) . Then

$$(b-x)^2 B - (x-a)^2 A = (B-A) \left[x - \left(\frac{bB-aA}{B-A} \right) \right]^2 - \frac{AB}{B-A} (b-a)^2$$

and by (2.11) we get

$$(2.18) \quad \begin{aligned} & (x-a)f(a) + (b-x)f(b) - \int_a^b f(t) dt \\ & \leq (B-A) \left[x - \left(\frac{bB-aA}{B-A} \right) \right]^2 - \frac{AB}{(B-A)^2} (b-a)^2 \end{aligned}$$

for any $x \in [a, b]$.

If $A \geq 0$, then $x_0 = \frac{bB-aA}{B-A} \in [a, b]$, and by (2.18) for $x = \frac{bB-aA}{B-A}$ we get that

$$(2.19) \quad 0 \leq \frac{1}{2} \cdot \frac{AB}{B-A} (b-a) \leq \frac{Bf(a) - Af(b)}{B-A} - \frac{1}{b-a} \int_a^b f(t) dt$$

which is an interesting inequality in itself as well.

3. THE COMPOSITE CASE

Consider the division $I_n : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ and denote $h_i := x_{i+1} - x_i$ ($i = \overline{0, n-1}$). If $\xi_i \in [x_i, x_{i+1}]$ ($i = \overline{0, n-1}$) are intermediate points, then we will denote by

$$(3.1) \quad G_n(f; I_n, \xi) := \sum_{i=0}^{n-1} [(\xi_i - x_i) f(x_i) + (x_{i+1} - \xi_i) f(x_{i+1})]$$

the generalised trapezoid rule associated to f , I_n and ξ .

The following theorem providing upper and lower bounds for the remainder in approximating the integral $\int_a^b f(t) dt$ of a convex function f in terms of the generalised trapezoid rule holds.

Theorem 7. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function and I_n and ξ be as above. Then we have:*

$$(3.2) \quad \int_a^b f(t) dt = G_n(f; I_n, \xi) - S_n(f; I_n, \xi),$$

where $G_n(f; I_n, \xi)$ is the generalised Trapezoid Rule defined by (3.1) and the remainder $S_n(f; I_n, \xi)$ satisfies the estimate:

$$\begin{aligned}
 (3.3) \quad & \frac{1}{2} \left[\sum_{i=0}^{n-1} (x_{i+1} - \xi_i)^2 f'_+ (\xi_i) - \sum_{i=0}^{n-1} (\xi_i - x_i)^2 f'_- (\xi_i) \right] \\
 & \leq S_n(f; I_n, \xi) \\
 & \leq \frac{1}{2} \left[(b - \xi_{n-1})^2 f'_- (b) + \sum_{i=1}^{n-1} \left[(x_i - \xi_{i-1})^2 f'_- (x_i) - (\xi_i - x_i)^2 f'_+ (x_i) \right] \right. \\
 & \quad \left. - (\xi_0 - a)^2 f'_+ (a) \right].
 \end{aligned}$$

Proof. If we write the inequalities (2.1) and (2.11) on the interval $[x_i, x_{i+1}]$ and for the intermediate points $\xi_i \in [x_i, x_{i+1}]$, then we have

$$\begin{aligned}
 & \frac{1}{2} \left[(x_{i+1} - \xi_i)^2 f'_+ (x_i) - (\xi_i - x_i)^2 f'_- (\xi_i) \right] \\
 & \leq (\xi_i - x_i) f(x_i) + (x_{i+1} - \xi_i) f(x_{i+1}) - \int_{x_i}^{x_{i+1}} f(t) dt \\
 & \leq \frac{1}{2} \left[(x_{i+1} - \xi_i)^2 f'_- (x_{i+1}) - (\xi_i - x_i)^2 f'_+ (x_i) \right].
 \end{aligned}$$

Summing the above inequalities over i from 0 to $n-1$, we deduce

$$\begin{aligned}
 (3.4) \quad & \frac{1}{2} \sum_{i=0}^{n-1} \left[(x_{i+1} - \xi_i)^2 f'_+ (\xi_i) - (\xi_i - x_i)^2 f'_- (\xi_i) \right] \\
 & \leq G_n(f; I_n, \xi) - \int_a^b f(t) dt \\
 & \leq \frac{1}{2} \left[\sum_{i=0}^{n-1} (x_{i+1} - \xi_i)^2 f'_- (x_{i+1}) - \sum_{i=0}^{n-1} (\xi_i - x_i)^2 f'_+ (x_i) \right].
 \end{aligned}$$

However,

$$\begin{aligned}
 \sum_{i=0}^{n-1} (x_{i+1} - \xi_i)^2 f'_- (x_{i+1}) &= (b - \xi_{n-1})^2 f'_- (b) + \sum_{i=0}^{n-2} \left[(x_{i+1} - \xi_i)^2 f'_- (x_{i+1}) \right] \\
 &= (b - \xi_{n-1})^2 f'_- (b) + \sum_{i=1}^{n-1} (x_i - \xi_{i-1})^2 f'_- (x_i)
 \end{aligned}$$

and

$$\sum_{i=0}^{n-1} (\xi_i - x_i)^2 f'_+ (x_i) = \sum_{i=1}^{n-1} (\xi_i - x_i)^2 f'_+ (x_i) + (\xi_0 - a)^2 f'_+ (a)$$

and then, by (3.4), we deduce the desired estimate (3.3). ■

The following corollary may be useful in practical applications.

Corollary 4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable convex function on $[a, b]$. Then we have the representation (3.2) and $S_n(f; I_n, \xi)$ satisfies the estimate:*

$$\begin{aligned}
 (3.5) \quad & \sum_{i=0}^{n-1} \left(\frac{x_i + x_{i+1}}{2} - \xi_i \right) h_i f'(\xi_i) \\
 & \leq S_n(f; I_n, \xi) \\
 & \leq \frac{1}{2} \left[(b - \xi_{n-1})^2 f'_-(b) - (\xi_0 - a)^2 f'_+(a) \right. \\
 & \quad \left. + \sum_{i=1}^{n-1} \left[\left(x_i - \frac{\xi_i + \xi_{i-1}}{2} \right) (\xi_i - \xi_{i-1}) f'(x_i) \right] \right].
 \end{aligned}$$

We may also consider the trapezoid quadrature rule:

$$(3.6) \quad T_n(f; I_n) := \sum_{i=0}^{n-1} \frac{f(x_i) + f(x_{i+1})}{2} \cdot h_i.$$

Using the above results, we may state the following corollary.

Corollary 5. *Assume that $f : [a, b] \rightarrow \mathbb{R}$ is a convex function on $[a, b]$ and I_n is a division as above. Then we have the representation*

$$(3.7) \quad \int_a^b f(t) dt = T_n(f; I_n) - Q_n(f; I_n)$$

where $T_n(f; I_n)$ is the mid-point quadrature formula given in (3.6) and the remainder $Q_n(f; I_n)$ satisfies the estimates

$$\begin{aligned}
 (3.8) \quad 0 & \leq \frac{1}{8} \sum_{i=0}^{n-1} \left[f'_+ \left(\frac{x_i + x_{i+1}}{2} \right) - f'_- \left(\frac{x_i + x_{i+1}}{2} \right) \right] h_i^2 \\
 & \leq Q_n(f; I_n) \leq \frac{1}{8} \sum_{i=0}^{n-1} [f'_+(x_{i+1}) - f'_-(x_i)] h_i^2.
 \end{aligned}$$

The constant $\frac{1}{8}$ is sharp in both inequalities.

4. APPLICATIONS FOR P.D.F.S

Let X be a random variable with the probability density function $f : [a, b] \subset \mathbb{R} \rightarrow [0, \infty)$ and with cumulative distribution function $F(x) = \Pr(X \leq x)$.

The following theorem holds.

Theorem 8. *If $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}_+$ is monotonically increasing on $[a, b]$, then we have the inequality:*

$$\begin{aligned}
 (4.1) \quad & \frac{1}{2} \left[(b - x)^2 f_+(x) - (x - a)^2 f_-(x) \right] + x \\
 & \leq E(X) \\
 & \leq \frac{1}{2} \left[(b - x)^2 f_+(b) - (x - a)^2 f_-(a) \right] + x
 \end{aligned}$$

for any $x \in (a, b)$, where $f_{\pm}(\alpha)$ represent respectively the right and left limits of f in α and $E(X)$ is the expectation of X .

The constant $\frac{1}{2}$ is sharp in both inequalities.

The second inequality also holds for $x = a$ or $x = b$.

Proof. Follows by Theorem 5 and 6 applied for the convex cdf function $F(x) = \int_a^x f(t) dt$, $x \in [a, b]$ and taking into account that

$$\int_a^b F(x) dx = b - E(X).$$

■

Finally, we may state the following corollary in estimating the expectation of X .

Corollary 6. *With the above assumptions, we have*

$$(4.2) \quad \begin{aligned} & \frac{1}{8} \left[f_+ \left(\frac{a+b}{2} \right) - f_- \left(\frac{a+b}{2} \right) \right] (b-a)^2 + \frac{a+b}{2} \\ & \leq E(X) \leq \frac{1}{8} [f_+(b) - f_-(a)] (b-a)^2 + \frac{a+b}{2}. \end{aligned}$$

5. APPLICATIONS FOR HH -DIVERGENCE

Assume that a set χ and the σ -finite measure μ are given. Consider the set of all probability densities on μ to be

$$(5.1) \quad \Omega := \left\{ p|p : \Omega \rightarrow \mathbb{R}, p(x) \geq 0, \int_{\chi} p(x) d\mu(x) = 1 \right\}.$$

Csiszár's f -divergence is defined as follows [4]

$$(5.2) \quad D_f(p, q) := \int_{\chi} p(x) f \left[\frac{q(x)}{p(x)} \right] d\mu(x), \quad p, q \in \Omega,$$

where f is convex on $(0, \infty)$. It is assumed that $f(u)$ is zero and strictly convex at $u = 1$. By appropriately defining this convex function, various divergences are derived.

In [5], Shioya and Da-te introduced the generalised Lin-Wong f -divergence $D_f(p, \frac{1}{2}p + \frac{1}{2}q)$ and the Hermite-Hadamard (HH) divergence

$$(5.3) \quad D_{HH}^f(p, q) := \int_{\chi} \frac{p^2(x)}{q(x) - p(x)} \left(\int_1^{\frac{q(x)}{p(x)}} f(t) dt \right) d\mu(x), \quad p, q \in \Omega,$$

and, by the use of the Hermite-Hadamard inequality for convex functions, proved the following basic inequality

$$(5.4) \quad D_f \left(p, \frac{1}{2}p + \frac{1}{2}q \right) \leq D_{HH}^f(p, q) \leq \frac{1}{2} D_f(p, q),$$

provided that f is convex and normalised, i.e., $f(1) = 0$.

The following result in estimating the difference

$$\frac{1}{2} D_f(p, q) - D_{HH}^f(p, q)$$

holds.

Theorem 9. Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a normalised convex function and $p, q \in \Omega$. Then we have the inequality:

$$\begin{aligned}
 (5.5) \quad 0 &\leq \frac{1}{8} \left[D_{f'_+ \cdot |\cdot|^{-\frac{1}{2}}} (p, q) - D_{f'_- \cdot |\cdot|^{-\frac{1}{2}}} (p, q) \right] \\
 &\leq \frac{1}{2} D_f (p, q) - D_{HH}^f (p, q) \\
 &\leq \frac{1}{8} D_{f'_- \cdot (-1)} (p, q).
 \end{aligned}$$

Proof. Using the double inequality

$$\begin{aligned}
 0 &\leq \frac{1}{8} \left[f'_+ \left(\frac{a+b}{2} \right) - f'_- \left(\frac{a+b}{2} \right) \right] |b-a| \\
 &\leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \\
 &\leq \frac{1}{8} [f_-(b) - f'_+(a)] (b-a)
 \end{aligned}$$

for the choices $a = 1$, $b = \frac{q(x)}{p(x)}$, $x \in \chi$, multiplying with $p(x) \geq 0$ and integrating over x on χ we get

$$\begin{aligned}
 0 &\leq \frac{1}{8} \int_{\chi} \left[f'_+ \left(\frac{p(x) + q(x)}{2p(x)} \right) - f'_- \left(\frac{p(x) + q(x)}{2p(x)} \right) \right] |q(x) - p(x)| d\mu(x) \\
 &\leq \frac{1}{2} D_f (p, q) - D_{HH}^f (p, q) \\
 &\leq \frac{1}{8} \int_{\chi} \left[f'_- \left(\frac{q(x)}{p(x)} \right) - f'_+ (1) \right] (q(x) - p(x)) d\mu(x),
 \end{aligned}$$

which is clearly equivalent to (5.5). ■

Corollary 7. With the above assumptions and if f is differentiable on $(0, \infty)$, then

$$(5.6) \quad 0 \leq \frac{1}{2} D_f (p, q) - D_{HH}^f (p, q) \leq \frac{1}{8} D_{f'_- \cdot (-1)} (p, q).$$

REFERENCES

- [1] P. CERONE and S.S. DRAGOMIR, Trapezoidal-type rules from an inequalities point of view, in *Handbook of Analytic-Computational Methods in Applied Mathematics*, G.A. Anastassiou (Ed), Chapman & Hall/CRC Press, New York, 2000, 65-134.
- [2] P. CERONE, S.S. DRAGOMIR and C.E.M. PEARCE, A generalised trapezoid inequality for functions of bounded variation, *Turkish J. of Math.*, **24**(2) (2000), 147-163.
- [3] S.S. DRAGOMIR, On the trapezoid quadrature formula for Lipschitzian mappings and applications, *Tamkang J. of Math.*, **30**(2) (1999), 133-138.
- [4] I. CSISZÁR, Information-type measures of difference of probability distributions and indirect observations, *Studia Math. Hungarica*, **2** (1967), 299-318.
- [5] H. SHIOYA and T. DA-TE, A generalisation of Lin divergence and the derivation of a new information divergence, *Elec. and Comm. in Japan*, **78**(7), (1995), 37-40.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO BOX 14428, MELBOURNE CITY MC, 8001, VICTORIA, AUSTRALIA.

E-mail address: sever@matilda.vu.edu.au

URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>